

[This question paper contains 4 printed pages.]

**Your Roll No.....**

**Sr. No. of Question Paper : 851**

**G**

**Unique Paper Code : 2352572301**

**Name of the Paper : Differential Equations**

**Name of the Course : B.Sc. (Physical Science and  
Mathematical Science) with  
Operational Research and  
Bachelor of Arts**

**Semester : III**

**Duration : 3 Hours**

**Maximum Marks : 90**

**Instructions for Candidates**

1. Write your Roll No. on the top immediately on receipt of this question paper.
2. Attempt all questions by selecting two parts form each question.
3. All questions carry equal marks.

1. (a) Find the general solution of the Bernoulli equation

given by  $\frac{dy}{dx} + \frac{y}{2x} = \frac{x}{y^3}$  with initial condition  $y(1) = 2$ .

Also find an integrating factor for the linear differential equation

$$\frac{dy}{dx} + \left( \frac{2x+1}{x} \right) y = e^{-2x}. \quad (7\frac{1}{2})$$

**P.T.O.**

- (b) Find the general solution of the differential equation  $(x^2 - 3y^2)dx + 2xy dy = 0$  by showing it's a homogeneous equation. Also show that  $M(tx, ty)$

$$= tM(x, y) \text{ for } M = y + \sqrt{x^2 + y^2}. \quad (7\frac{1}{2})$$

- (c) Determine the most general  $N(x, y)$  for the equation  $(x^{-2}y^{-2} + xy^{-3})dx + N(x, y)dy = 0$  such that the equation is exact and solve the resulting exact equation. (7½)

2. (a) Assume that the population of a certain city increase at a rate proportional to the number of inhabitants at any time, if the population doubles in 40 years, in how many years will it triple?

(7½)

- (b) Show that the relation  $x^2 + y^2 - 25 = 0$  is an implicit solution of the differential equation

$x + y \frac{dy}{dx} = 0$  on the interval  $-5 < x < 5$ . Explain whether the relation  $x^2 + y^2 + 25$  is also an implicit solution of  $x + y \frac{dy}{dx} = 0$ . (7½)

- (c) Find the particular solution of the linear system that satisfies the stated initial conditions :

$$\frac{dy_1}{dt} = y_1 + y_2, \quad y_1(0) = 1$$

$$\frac{dy_2}{dt} = 4y_1 + y_2, \quad y_2(0) = 6. \quad (7\frac{1}{2})$$

3. (a) Solve the initial value problem :

$$x^2 y'' + 3xy' + y = 0, \quad y(1) = 4, \quad y'(1) = -1. \quad (7\frac{1}{2})$$

- (b) Find a homogeneous linear ordinary differential equation for which two functions  $x^{-3}$  and  $x^{-3} \ln x$  ( $x > 0$ ) are solutions. Show also linear independence by considering their Wronskian.

(7½)

- (c) Consider the initial value problem :

$$\frac{dy}{dx} = 1 + y^2, \quad y(0) = 0.$$

Examine the existence and uniqueness of solution in the rectangle :  $|x| < 5, |y| < 3$ . (7½)

4. (a) Find a general solution of the following nonhomogeneous differential equation :

$$y'' + 4y' + 4y = e^{-2x} \sin 2x. \quad (7\frac{1}{2})$$

- (b) Use the method of undetermined coefficients to find the particular solution of the differential equation :

$$y'' - 2y' + y = x^2 + e^x. \quad (7\frac{1}{2})$$



- (c) Use the method of variation of parameters to find a particular solution of the differential equation :

$$y'' + y = \tan x \sec x. \quad (7\frac{1}{2})$$

5. (a) Find the general solution of the equation.

$$(x - y)y^2u_x + (x - y)x^2u_y = (x^2 + y^2)u \quad (7\frac{1}{2})$$

- (b) Eliminate the constants  $a$  and  $b$  from the equation

$$2z = (ax + y)^2 + b \quad (7\frac{1}{2})$$

- (c) Solve the initial value problem :

$$u_t + uu_t = x, \quad u(x, 0) = 1 \quad (7\frac{1}{2})$$

6. (a) Find the general solution of the linear partial differential equation.

$$x(y^2 - z^2)u_x + y(z^2 - x^2)u_y + z(x^2 - y^2)u_z = 0 \quad (7\frac{1}{2})$$

- (b) Use  $v = \ln u$  and  $v = f(x) + g(y)$  to solve the equation.

$$x^2 u_x^2 + y^2 u_y^2 = u^2 \quad (7\frac{1}{2})$$

- (c) Reduce the equation:  $x^2 u_{xx} + 2x u_{xy} + y^2 u_{yy} = 0$  to canonical form and hence find the general solution. (7\frac{1}{2})