[This question paper contains 8 printed pages.]

Your Roll No.....

Sr. No. of Question Paper: A

Unique Paper Code 32351402

: BMATH-409; Riemann Name of the Paper

Integration and Series of

Functions

: B.Sc. (H) Mathematics Name of the Course

: IV Semester

Maximum Marks: 75 Duration: 3 hours + 30 minutes

Instructions for Candidates

- Write your Roll No. on the top immediately on receipt 1. of this question paper.
- Attempt any two parts from each question. 2.
- All questions are compulsory. 3.
- (a) Let f be a bounded function on [a, b]. Define 1. integrability of f on [a, b] in the sense of Riemann.

- (b) Prove that every continuous function on [a, b] is integrable. Discuss about the integrability of discontinuous functions.
- (c) Let $f(x) = \sin \frac{1}{x}$ for $x \neq 0$ and f(0) = 0. Show that $|\int_{-1}^{1} f(t) dt| \leq 2.$ (6)
- (d) Let f(x) = x for rational x; and f(x) = 0 for irrational x. Calculate the upper and lower Darboux integrals of f on the interval [0, b]. Is f integrable on [0, b]?
- 2. (a) State Fundamental Theorem of Calculus II. Use it to calculate

$$\lim_{x \to 0} \frac{1}{x} \int_0^x e^{t^2} dt.$$
 (6.5)

(b) Let f be defined as (6.5)

$$f(t) = \begin{cases} 0, & t < 0 \\ t, & 0 \le t \le 1 \\ 4, & t > 1 \end{cases}$$

- (i) Determine the function $F(x) = \int_0^x f(t) dt$.
- (ii) Sketch F. Where is F continuous?
- (iii) Where is F differentiable? Calculate F' at points of differentiability.
- (c) State and prove Intermediate value theorem for Integral Calculus. Give an example to show that condition of continuity of the function cannot be relaxed. (6.5)
- (d) Let f be a continuous function on R. Define

$$G(x) = \int_0^{\sin x} f(t)dt$$
 for $x \in \mathbb{R}$.

Show that G is differentiable on \mathbb{R} and compute G'. (6.5)

3. (a) Let $\beta(p, q)$ (where p, q > 0) denotes the beta function, show that

$$\beta(p,q) = \int_{0^{+}}^{\infty} \frac{u^{p-1}}{(1+u)^{p+q}} du = \int_{0^{+}}^{1} \frac{v^{p-1} + v^{q-1}}{(1+v)^{p+q}} dv.$$
 (6)

(b) Determine the convergence and divergence of the following improper integrals

(i)
$$\int_0^1 \frac{\mathrm{d}x}{x(\ln x)^2}$$

(ii)
$$\int_{-1}^{\infty} \frac{dx}{x^2 + 4x + 6}$$
 (6)

(c) Define Improper Integral of type II.

Show that the improper integral $\int_{1}^{\infty} \frac{dx}{x^{p}}$ converges iff p > 1.

- (d) Show that the improper integral $\int_{\pi}^{\infty} \frac{\sin x}{x} dx$ is convergent but doesn't converge absolutely.
- 4. (a) Let $\langle f_n \rangle$ be a sequence of integrable functions on [a,b] and suppose that $\langle f_n \rangle$ converges uniformly on [a, b] to f. Show that is f is integrable.

- (b) Define
 - (i) pointwise convergence of sequence of functions
 - (ii) uniform convergence of a sequence of functions
 - (iii) If $A \subseteq \mathbb{R}$ and $\emptyset: A \to \mathbb{R}$ then define uniform norm of \emptyset on A. (6)
- (c) (i) Discuss the pointwise and uniform $\text{convergence of } f_n(x) = \frac{x}{n} \text{ for } x \in \mathbb{R}, \, n \in \mathbb{N}.$
 - (ii) Show that the sequence $\langle f_n \rangle$ where $f_n(x) = \frac{n}{x+n}$, $x \ge 0$ is uniformly convergent in any finite interval. (6)
- (d) (i) Show that the sequence $< f_n >$ where $f_n(x) = \frac{\sin nx}{\sqrt{n}}$ uniformly convergent on $[0, \pi]$.
 - (ii) Discuss the pointwise and uniform convergence of the sequence $g_n(x) = x^n$ for $x \in \mathbb{R}$, $n \in \mathbb{N}$.

- 5. (a) (i) State and prove the Cauchy Criteria for uniform convergence of series. (3.5)
 - (ii) Show that the series $\sum_{0}^{\infty} (1-x)x^{n}$ is not uniformly convergent on [0, 1] (3)
 - (b) Show that $\sum \frac{(-1)^n}{n^p} \frac{x^{2n}}{(1+x^{2n})}$ converges absolutely and uniformly for all values of x if p > 1.
 - (c) Is the sequence $< f_n >$ where $f_n = \frac{\sin(nx+n)}{n}$, uniformly convergent on \mathbb{R} ? Justify. (6.5)
 - (d) If f_n is continuous on $D \subseteq \mathbb{R}$ to \mathbb{R} for each $n \in N$ and $\sum f_n$ converges to f uniformly on D then prove that f is continuous on D. (6.5)
- 6. (a) Find the radius of convergence and exact interval of convergence of the following power series:

(i)
$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2} (x+1)^n$$

(ii)
$$\sum_{n=0}^{\infty} x^{n!}$$
 (6.5)

(b) Let $f(x) = \sum_{n=0}^{\infty} a_n x^n$ has radius of convergence R > 0. Show that the function f is differentiable on (-R, R) and

$$f'(x) = \sum_{n=1}^{\infty} na_n x^{n-1} \text{ for } |x| < R.$$
 (6.5)

(c) Show that

(i)
$$\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + ---$$

for $|x| < 1$

(ii)
$$\log 2 = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - - - - -$$
 (6.5)

(d) Let $f(x) = \sum_{n=0}^{\infty} a_n x^n$ be a power series with radius of convergence R > 0. If $0 < R_1 < R$, show that the power series converges uniformly on $[-R_1, R_1]$. Also, show that the sum function f(x) is continuous on the interval (-R, R). (6.5)

(3100)