[This question paper contains 4 printed pages.]

Your Roll No.....

Sr. No. of Question Paper: 1944

A

Unique Paper Code : 32355444

Name of the Paper : Elements of Analysis

Name of the Course : Mathematics: Generic

Elective for Honours

Semester : IV

Duration: 3 Hours Maximum Marks: 75

## Instructions for Candidates

- 1. Write your Roll No. on the top immediately on receipt of this question paper.
- 2. All questions are compulsory.
- 3. Attempt any two parts from each question.
- 1. a) Define countable and uncountable sets. Show that the set  $\mathbb{Q}$  of all rational numbers is denumerable. (7.5)
  - b) (i) Show that if  $a, b \in \mathbb{R}$ ,  $\max\{a, b\} = \frac{1}{2}(a + b + |a b|)$  and  $\min\{a, b\} = \frac{1}{2}(a + b |a b|)$ .
    - (ii) Determine the set A of all real numbers x such that |x 1| < |x|. (3.5+4)

- c) Let S be a bounded set in  $\mathbb{R}$  and let  $S_0$  be a nonempty subset of S. Show that  $\inf S \leq \inf S_0 \leq \sup S_0 \leq \sup S$ . (7.5)
- d) Using Archimedean property, show that  $\sup \left\{ 1 \frac{1}{n} : n \in \mathbb{N} \right\} = 1.$  (7.5)
- 2.a) Define a convergent sequence. Using the definition, show that  $\lim_{n\to\infty} (1 + \frac{(-1)^n}{n}) = 1$ . (7.5)
  - b) State and prove the squeeze theorem for sequences. Hence, show that  $\lim_{n\to\infty} \frac{\sin n}{n} = 0$ . (7.5)
  - c) (i) State Bolzano Weierstrass Theorem for sequences. Give an example of an unbounded sequence that has exactly two limit points.
    - (ii) Define limit superior and limit inferior of a sequence. For  $\langle x_n \rangle = \langle (-1)^n + \frac{2}{n} \rangle$ , show that  $\lim_{n \to \infty} \sup(-x_n) = -\lim_{n \to \infty} \inf(x_n)$ . (2+5.5)
- d) Define Monotone Sequences. Let the sequence  $\langle x_n \rangle$  be defined by  $x_1 = 1$  and  $x_{n+1} = 2 \frac{1}{x_n}$  for  $n \in \mathbb{N}$ . Show that the sequence  $\langle x_n \rangle$  is monotone and bounded.

Find its limit. (7.5)

- 3.a) State the Integral test. Show using integral test that the series  $\sum_{n=2}^{\infty} \frac{1}{n(\ln n)}$  is convergent. (6.5)
- b) Check whether the series are convergent or divergent:

i) 
$$\sum_{n=1}^{\infty} \frac{1}{3^{n}+1}.$$
ii) 
$$\sum_{n=1}^{\infty} \frac{n^{2}-1}{n^{4}}.$$
 (3+3.5)

c) State D'Alembert Ratio test. Construct examples of p-series where the ratio test fails.

(6.5)

- d) State Absolute Convergence test and Conditional Convergence test. Show that the converse of absolute convergence test is not true. (6.5)
- 4. a) Test the series  $\sum_{n=1}^{\infty} \frac{(n+1)^{n^2}}{(n)^{n^2}}$  for convergence. (6)
  - b) Determine whether the following series converge or diverge:

$$i)\sum\nolimits_{n=1}^{\infty}\frac{1}{2n-1}.$$

ii) 
$$\sum_{n=1}^{\infty} \frac{(n!)^2}{(2n)!}$$
. (3+3)

- c) Check the convergence of the series  $\sum_{n=1}^{\infty} \frac{\sqrt{n+1} \sqrt{n-1}}{n}.$  (6)
- d) Examine the series  $\sum_{n=1}^{\infty} \frac{3^{n}+4^{n}}{5^{n}}$  for convergence. Find its sum. (6)
- 5.a) Define Radius of convergence of a power series. Give examples of the series having radius of convergence 0 and ∞. (5)
  - b) Find interval of convergence for the power series  $\sum_{n=1}^{\infty} \frac{(-1)^n x^n}{n}.$  (5)
  - c) Using Differentiation theorem for power series, show that

$$\sum_{n=1}^{\infty} n^2 x^n = \frac{x(1+x)}{(1-x)^3} \tag{5}$$

d) Define cosine and sine functions as sums of power series. Show that for every  $x, y \in \mathbb{R}$ .

(i) 
$$S(x-y) = S(x) C(y) - C(x)S(y)$$

(ii) 
$$S'(x) = C(x)$$
 and  $S(2x) = 2S(x)C(x)$ . (2.5+2.5)

- 6.a) Find the interval of convergence of the power series  $\sum_{n=1}^{\infty} \frac{(x-2)^n}{n^2}.$  (5)
  - b) Using the power series of the function  $f(x) = \frac{1}{(1-x)}$  valid for |x| < 1 show that  $\frac{1}{x} = \sum_{n=0}^{\infty} (-1)^n (x-1)^n.$  (5)
- c) State Integration theorem for power series For |x| < 1, find a power series representation for  $\frac{1}{1+x^2}$ . Show that  $\tan^{-1}(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1}$ . (5)
- d) Write the power series expansion for  $f(x) = e^x$  about  $x_0 = 0$ . (5)