

[This question paper contains 4 printed pages.]

Your Roll No.....

Sr. No. of Question Paper : 4997

E

Unique Paper Code : 62351201

Name of the Paper : Algebra

Name of the Course : B.A. (Prog.)

Semester : II

Duration : 3 Hours

Maximum Marks : 75

**Instructions for Candidates**

1. Write your Roll No. on the top immediately on receipt of this question paper.
2. Attempt **all** questions.
3. Attempt any **two** parts from each question.
4. Marks are indicated against each question.

1. (a) Form an equation whose roots are  $-1, 2, 3 \pm 2i$ .

(6)

(b) Solve the equation

(6)

$$x^3 - 13x^2 + 15x + 189 = 0,$$

being given that one of the roots exceeds another by 2.

(c) If  $\alpha, \beta, \gamma$ , be the roots of the equation

(6)

$x^3 + 5x^2 - 6x + 3 = 0$ , find the value of

P.T.O.

$$(i) \sum \alpha^3 \qquad (ii) \sum (\alpha - \beta)^2 .$$

2. (a) Prove that : (6.5)

$$2^{10} \cos^6 \theta \sin^5 \theta = \sin 11\theta + \sin 9\theta - 5 \sin 7\theta - 5 \sin 5\theta + 10 \sin 3\theta + 10 \sin \theta.$$

(b) Sum the series : (6.5)

$$\cos \alpha + \cos (\alpha + \beta) + \cos (\alpha + 2\beta) + \dots \text{ to } n \text{ terms, provided } \beta \neq 2k\pi.$$

(c) State DeMoivre's theorem for rational indices and use it to solve the equation : (6.5)

$$x^7 - x^4 + x^3 - 1 = 0.$$

3. (a) Find the characteristic roots of the matrix A where (6)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & -4 & 2 \\ 0 & 0 & 7 \end{bmatrix}$$

(b) Solve the system of linear equations (6)

$$2x - 5y + 7z = 6$$

$$x - 3y + 4z = 3$$

$$3x - 8y + 11z = 11$$

- (c) Using Cayley Hamilton's Theorem, find the value of  $A^3$ , where (6)

$$A = \begin{bmatrix} -1 & 1 & 2 \\ 0 & 1 & -1 \\ 2 & 2 & 1 \end{bmatrix}$$

4. (a) Let  $X$  and  $Y$  be two subspace of a vector space  $V$ . (6.5)

(i) Prove that the intersection  $X \cap Y$  is also subspace of  $V$ .

(ii) Show that the union  $X \cup Y$  need not be a subspace of  $V$ .

- (b) Let  $V = F[a, b]$  be the set of all real valued functions defined on the interval  $[a, b]$ . For any  $f$  and  $g$  in  $V$ ,  $c$  in  $R$ , we define

$$(f + g)(x) = f(x) + g(x),$$

$$(c.f)(x) = cf(x)$$

Prove that  $V$  is a vector space over  $R$ , where  $R$  denotes the set of real numbers. (6.5)

- (c) Show that the vectors  $v_1 = (1,1,1)$ ,  $v_2 = (1,1,0)$ ,  $v_3 = (1,0,0)$  form a spanning set of  $R^3(R)$ , where  $R$  denotes the set of real numbers. (6.5)

P.T.O.

5. (a) Find the multiplicative inverse of the given elements (if it exists) if it does not exist, give the reason

(i)  $[12]$  in  $Z_{16}$       (ii)  $[38]$  in  $Z_{83}$       (6)

- (b) Find the order of each of the following permutations

(i)  $f = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 6 & 4 & 5 & 3 & 2 & 1 \end{pmatrix}$

(ii)  $f = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 4 & 6 & 7 & 5 & 1 & 8 & 2 & 3 \end{pmatrix}$

- (c) Let  $G$  be a group. Prove that  $G$  is abelian if and only if  $(ab)^{-1} = a^{-1}b^{-1}$  for all  $a, b \in G$ .      (6)

6. (a) Prove that the set  $S = \{0, 2, 4, 6, 8\}$  is an abelian group with respect to addition modulo 10.      (6.5)

- (b) Let  $G$  be the group of all  $2 \times 2$  invertible matrices with real entries under the usual matrix multiplication. Show that subset  $S$  of  $G$  defined by

$$S = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix}, b=c \right\}, \text{ does not form a subgroup of } G. \quad (6.5)$$

- (c) Show that  $Q(\sqrt{2}) = \{a + b\sqrt{2} : a, b \in Q\}$  is a subring of  $R$ , where  $R$  is a set of real numbers &  $Q$  is set of rational numbers.      (6.5)